



(11) **EP 1 829 447 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:
15.08.2012 Bulletin 2012/33

(21) Application number: **05822470.0**

(22) Date of filing: **13.12.2005**

(51) Int Cl.:
A01N 3/00 (2006.01) A01N 49/00 (2006.01)

(86) International application number:
PCT/MX2005/000115

(87) International publication number:
WO 2006/078152 (27.07.2006 Gazette 2006/30)

(54) **AQUEOUS COMPOSITION FOR THE PROTECTION AND STRENGTHENING OF PLANTS AND APPLICATION METHOD THEREOF**

WÄSSRIGE ZUSAMMENSETZUNG FÜR DEN SCHUTZ UND DIE STÄRKUNG VON PFLANZEN, SOWIE ANWENDUNGSVERFAHREN HIERFÜR

COMPOSITION AQUEUSE DESTINEE A PROTEGER ET A FORTIFIER DES PLANTES ET PROCEDE D'APPLICATION CORRESPONDANT

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI SK TR

(30) Priority: **17.12.2004 MX JL04000049**

(43) Date of publication of application:
05.09.2007 Bulletin 2007/36

(73) Proprietor: **Centro de Investigación y Asistencia en Tecnología y Diseño del Estado de Jalisco, A.C. (CIATEJ, A.C.)**
Guadalajara,
Jalisco 44270 (MX)

(72) Inventors:
• **RODRÍGUEZ GARAY, Benjamín**
Col. Las Fuentes CP 45070 (MX)
• **URBINA LÓPEZ, Dionicio**
C.P. 44300 Guadalajara, Jalisco (MX)
• **RODRÍGUEZ DOMÍNGUEZ, José Manuel**
C.P. 44270 Guadalajara, Jalisco (MX)

(74) Representative: **ABG Patentes, S.L.**
Avenida de Burgos, 16D
Edificio Euromor
28036 Madrid (ES)

(56) References cited:
WO-A1-91/03159 WO-A1-96/24246
WO-A2-2004/095926 GB-A- 2 287 637
SU-A1- 1 153 403 US-A- 4 710 394
US-A- 4 828 890

EP 1 829 447 B1

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

TECHNICAL FIELD

[0001] The present invention relates to the techniques used in the agriculture industry and gardening for the obtainment and manufacturing of compositions useful in cultures, management and caring of plants, and more particularly, it is related to an aqueous composition for the protection and strengthening of plants as well as to the application method thereof.

BACKGROUND OF THE INVENTION

[0002] As it is well known, plants suffer of several types of biotic or abiotic stress reducing their vitality and some times provoking the death of the plants.

[0003] For example, in the micro-propagation biotechnological processes, the plants become dehydrated and photo-oxidized due to a sudden light excess and intensity received by the plant when from an *in vitro* condition goes through an adaptation phase to the *ex vitro* life, i.e., in the micro-propagation, the plants suffer a sudden change of light conditions, and in the *ex vitro* phase are positioned in a soil located in any type of container. In addition, through all the *ex vitro* adaptation phase and up to the plants being capable to their transplanting in the final destination, the plants are exposed to stress conditions. Moreover, in said transplanting the plants suffer a very important dehydrating stress due to the sudden changes in going to a lower relative humidity conditions compared to the inside *in vitro* conditions.

[0004] Similarly, the same problem is observed in the small leaf clusters obtained by conventional germination systems or by cuttings obtained in greenhouses. In this regard, a good health plant has a higher commercial value than one stressed, being this an important factor for the survival of the field vegetables.

[0005] Regarding the dehydration, it is worth mentioning that this phenomenon may also be caused by a lack of watering or by the high speed winds, which make the plants to loss water through the leaves. Moreover, it is worth to remember that the plants transpire water through their leaves due to the photosynthetic activity and breathing process.

[0006] All the above results in the existence of damaged plants requiring compounds or substances that not only help to prevent dehydration conditions, but in addition help the plants to recover from the bath health in a few time, since in case of a non opportune action, the plants may become more deteriorated.

[0007] On the other hand, the plants also suffer diseases and are attacked by pests, such as the white fly being a very common pest in greenhouse cultures and even in the field, further, the white fly transmits viral infections damaging the plants in great extent.

[0008] To all the above mentioned factors causing plant damage or stress, the damage caused by man may

be added, when he does not properly manage the plants in the greenhouses or in the culture fields.

[0009] The above problems results in the prior art development of compositions trying to protect the plants against some of these factors causing stress, such as the composition described in the United States Patent No. 4,094,845, which is directed to provide a water loss reducing composition in the existing leaves and stems of plants by transpiration, besides, it has a protection effect against damages caused by the wind and cold. More specifically, the composition consists essentially of an aqueous solution or dispersion of a carboxylated hydrophilic acrylic polymer, a cross-linking agent for said polymer and an effective amount of UV rays absorption agent. The main feature of this composition is that have effect only to avoid water loosing, then having a limited application. In addition, it does not mention if a dehydrated plant can recover from this bad state.

[0010] Document WO96/24246 A1 discloses an aqueous film-forming composition containing a film-forming substance, and anti-oxidant e.g. beta carotene and a colouring agent e.g. a red colouring agent for coating seeds to protect them from ageing damage.

[0011] Document WO91/003159 describes the addition of polyols, wherein in a preferred embodiment uses several combinations such as glycerol and sorbitol; sorbitol and polyethylene glycol (PEG) and sorbitol, mannitol and polyethylene glycol. It is important to establish that the essential polyols are sorbitol and mannitol, and that PEG is not essential for the aim of said invention. Else, the salts used in the formulation are in fact the nutrient medium for the composition, not the other elements.

[0012] Document US 4,710,394 A discloses an aqueous solution for the preservation of plants containing citric acid, a dye and an inorganic salt.

[0013] Document SU 1,153,403 A1 discloses a composition for protecting plants from dehydration containing a polyethylene wax, a photo sensitiver and a green anthraquinone dye.

[0014] Document US 4,828,890 A discloses a method of preserving plants by perfusing into living plants and aqueous preservative compositions made of humectants, thus reducing bleeding of the preservative material.

[0015] As may be seen from the above, there is currently a lack of compositions which, with a few components, be highly effective for the protection of plants against several biotic or abiotic factors generating stress in plants, and above all, there is no composition which immediately improves, in a few time, the plants health.

In this regard, it is known that any kind of fertilizers help to improve the plants health but with a very low action, such that the benefits from the fertilizers cannot be perceived immediately. Besides, most of the compositions have an agricultural main use, leaving aside people liking gardening.

SUMMARY OF THE INVENTION

[0016] As a consequence from the above, it has been looking for suppressing the existing problems for the protection and strengthening of plants, in such a manner that it has been found that a composition comprising polyethylene glycol, vitamin A or a derivative or precursor thereof, including mixtures of same, and at least a red coloring agent, all components having a synergistic effect providing a protection and strengthening effect on plants, pointing out particularly a readily and surprisingly strengthening of the plants having stress conditions. Further, the compositions protect the plants from the stress through their growing and development. More precisely the aqueous composition for the protection and strengthening of plants of the present invention comprises

- a) polyethylene glycol in a concentration of about 2.0 to about 10.0 grams per liter of the composition, wherein the polyethylene glycol has a molecular weight of about 6000 to about 8000 grams/mol;
- b) vitamin A, or a derivative or precursor thereof, including mixtures of same, in a concentration of about 0.1 to about 10.0 grams per liter of the composition, and
- c) at least a red coloring agent in a concentration of about 0.1 to about 10.0 grams per liter of the composition, wherein said coloring agent enables the passage of light at wavelengths greater than 600 nanometers;

wherein the vitamin A derivative is selected from the group consisting of vitamin A acetate and vitamin A palmitate, and wherein said vitamin A precursor is selected from the group consisting of beta-carotene, zeaxanthine and luteine.

[0017] The red coloring agent is preferably a food grade coloring agent.

[0018] In a first embodiment of the present invention, the aqueous composition for the protection and strengthening of plants further comprises: (d) at least a glucoside in a concentration of about 0.5 to about 15 milligrams per liter of the composition; and/or (e) at least a foliar fertilizer.

[0019] Regarding the composition application method of the present invention, it comprises at least it: (a) providing a plant; and (b) applying the composition onto the leaves of the plant in an application dose of about 0.03 to about 0.05 ml per cm² of leaf of the plant, preferably, the application is made onto the upper side of the leaf or bundle receiving the sun light. As a preferred embodiment of the method of the present invention, the application is carried out by mechanical or manual spraying.

[0020] The composition of the present invention has outstanding effects when applied to plants obtained by micro-propagation going from *in vitro* conditions to *ex vitro* conditions, which are very susceptible to suffer stress, however, it can also be applied to plants obtained by conventional methods used in greenhouses.

[0021] In addition, an object of the present invention is to provide a protection and strengthening composition for plants, easy to obtain and to apply, and further having a lower cost compared with current compositions and fertilizers. A further object of the present invention is to provide a composition for the protection and strengthening of plants being innocuous for the man and other animal species.

BRIEF DESCRIPTION OF THE DRAWINGS

[0022] The novel aspects considered characteristics of the present invention are set forth particularly in the appended claims. However, the invention itself, both for its organization and for its application method, together with other objects and advantages thereof, will be better understood in the following detailed description of certain specific embodiments, when read in relation with the accompanying drawings, wherein:

Figure 1A is a side, bottom, perspective view of a plant of the species *Paulownia elongata* showing withered leaves due to a bad management in the greenhouse.

Figure 1B is a bottom, side, perspective view of the same plant showed in Figure 1A after 60 minutes from the receipt of the application of the aqueous composition of the present invention formulated according to a specific embodiment thereof.

Figure 2 is a comparative view showing two small leaf clusters of the "saladet" variety, cultured under the same greenhouse conditions, one of them having being treated with the composition of the present invention.

Figure 3 is a graphic showing the results obtained from a multiple media analysis, from height measurements carried out over study groups formed by small leaf clusters of "saladet" tomato obtained in a greenhouse, to which, except to one group, the composition of the present invention formulated according to specific embodiments thereof was applied.

Figure 4 is a graphic showing the results obtained in a multiple media analysis from height measurements carried out over study groups formed by small leaf clusters of "saladet" tomato transplanted to field, to which except to one group, the composition of the present invention, formulated according to specific embodiments thereof was applied.

Figure 5 is a graphic showing the results obtained in a multiple media analysis from measurements of the stem base diameter conducted over the same study groups of Figure 4.

DETAILED DESCRIPTION OF THE INVENTION

[0023] It has been found that an aqueous composition comprising: polyethylene glycol, vitamin A or a derivative or precursor thereof, including mixtures thereof, and at

least a red coloring agent, has a synergistic effect over its components that, when applied onto the plants, brings protection against factors causing stress to the plants that are in this state. The aqueous composition of the present invention more particularly comprises the following:

- a) polyethylene glycol in a concentration from about 2.0 to about 10.0 grams per liter of the composition, wherein the polyethylene glycol has a molecular weight of about 6000 to about 8000 grams/mol.
- b) vitamin A or a derivative or precursor thereof, including mixtures of same, in a concentration of about 0.1 to about 10.0 grams per liter of the composition; and,
- c) at least a red coloring agent in a concentration of about 0.1 to about 10.0 grams per liter of the composition, wherein said coloring agent enables the passage of the light at wavelengths greater than 600 nanometers;

wherein the vitamin A derivative is selected from the group consisting of vitamin A acetate and vitamin A palmitate, and wherein said vitamin A precursor is selected from the group consisting of beta-carotene, zeaxanthine and luteine.

[0024] With regard to the above, it can be mentioned that, with respect to the polyethylene glycol molecular weight, lower than the limits of the indicated ratio, the weight concentration of this component would be necessarily higher, it is to say, a great weight concentration of this component would be necessarily higher, it is to say, a great amount of this component would have to be used in the composition. On the other hand, if the polyethylene glycol molecular weight is higher than the ratio upper limit, it is very difficult to dissolve it in the aqueous media.

[0025] It is convenient to mention that the polyethylene glycol function within the composition is to act as adherent of the same over the leaves of the plants onto which it is applied, besides, the polyethylene glycol forms a transparent film reducing the water loss of the leaf by transpiration, as result of the plant photosynthesis and breathing.

[0026] Vitamin A and derivatives and precursors thereof have the function of avoiding the photo-oxidation and to counteract the noxious dehydration effect caused by the lack or loss of water, winds at high speed or other factors also causing this problem, further giving the plant heat resistance.

[0027] Regarding the red coloring agent, the same is preferably selected among food grade coloring agents with the purpose that the composition be easily manufactured and not to represent prejudicial effects to animal species, including humans. In a preferred manner, in the aqueous composition, a red coloring agent 3, a red coloring agent 40, or a mixture thereof is used. The main function of the red coloring agent is to form a film on the leaf surface which enables the passage of light at wave-

lengths greater than about 600 nm, thus avoiding the wavelengths corresponding to UV rays, which, as it is well known, cause great damage to living beings.

[0028] Describing a first embodiment of the present invention, the protection and strengthening aqueous solution for plants further comprises; (d) at least a glucoside in a concentration of about 0.5 to about 15 mg per liter of the composition, and/or, (e) at least a foliar fertilizer. From these, the glucoside functions as a natural agent against pathogen microorganisms and insects, besides to function as a stimulant for lignin production, being a compound giving consistency and strengthening to the plant cells, and as a consequence to the tissues, in a preferred manner said glucoside is selected from the following glucosides: phloridzin (phloretin-2-β-D-glucoside) phloretin (β-(4-hydroxyphenyl) 2,4,6-trihydroxypropionophenone), quercetin (3,3',4',5,6-pentahydroxyflavone), rutoside (quercetin-3-rutinoside) or phloroglucinol (1,3,5-trihydroxybenzene), in a preferred manner, the glucoside used in the invention is phloridzin.

[0029] Regarding the foliar fertilizers, these are widely used in the technical field, and among its main ingredients are the nitrogen, phosphorous and potassium compounds enriched with minerals such as chrome, iron, sulfur, zinc, calcium, magnesium; examples of these foliar fertilizers are those known under the commercial trademarks COSMOCEL 20-30-10®, TRICEL-20® and Bayfolan Forte®, these fertilizers are added to the aqueous composition according to the respective manufacturers recommended doses, it is worth to insist in the fact that the addition of this kind of fertilizers to the composition of the present invention is optional, since as it may be observed in the examples, the same has benefic effects either without the inclusion of these foliar fertilizers.

[0030] The pH of the aqueous composition for the protection and strengthening of plants of the present invention is between about 3.5 to about 7.0.

[0031] Regarding the application method for the aqueous composition for the protection and strengthening of plants of the present invention, said method comprises the following steps:

- a) providing a plant; and,
- b) applying the composition onto the leaves of the plant in an application dose of about 0.03 to about 0.05 ml per cm² of plant leaf.

[0032] With regard to the above, in a preferred embodiment of this method, the application is made over the upper side of the leaf (bundle) receiving the sun light or other source of light. The application dose in the indicated ratio permits to cover satisfactorily the whole leaf surface onto which the composition is being applied, out of the pointed ratio, the leaf would not be satisfactorily covered, or the composition would be excessively applied.

[0033] In another application method embodiment of the present invention, the same is realized by mechanical or manual spraying such that the application may be eas-

ily made by the greenhouse or culture field personnel, or by people having liking for gardening.

[0034] In a more preferred manner, the aqueous composition of the present invention is applied to plants obtained by micro-propagation going from *in vitro* conditions to *ex vitro* conditions, wherein it is known by the skilled in the art that plants suffer stress by a sudden greater light intensity receipt and by a sudden decrease of the environment relative humidity. However, the composition of the present invention may also be applied to small leaf clusters obtained by conventional seeds and cuttings methods, without being important the plant gender or species, such as vegetables, ornamental plants, etc.

[0035] The application of the present invention composition can be carried out at any time of the plant life, and the application may be repeated as required thereby.

[0036] The aqueous composition for the protection and strengthening of plants, as well as its application method will be more clearly illustrated by the following specific examples described below, which are only illustrative but not limitative of the invention, being said examples as follows:

EXAMPLE 1

Preparation of the composition

[0037] In a 1 L beaker with 500 ml bi-distillated water, 5 g of polyethylene glycol were dissolved under constant stirring, the molecular weight of the polyethylene glycol was 6000 grams/mol, further the following compounds were added one by one to the beaker: 0.5 g of pharmaceutical grade vitamin A palmitate for human use; and 0.5 g of red coloring agent under the trademark "Masterchef", which is a food grade coloring agent used in bakery, consisting of a mixture of red coloring agent 3 and red coloring agent 40. Finally, a sufficient amount of water was added to the beaker to complete a liter.

EXAMPLE 2

Application to bad managed stressed plants

[0038] In order to explain this example, reference to Figures 1A and 1B is made. Specifically, Figure 1A shows a plant 10 of the species *Paulownia elongata* of 4 weeks of life, the plant showed stress due to a bad management in the greenhouse, as a notorious evidence of the plant stress one of its leafs was completely withered, this leaf is clearly identified in Figure 1A within a circle and the 11 numeric reference.

[0039] With the object of improving the plant condition, the composition obtained in the example 1 was applied to the upper side of all the leafs of the plant, with an application dose of 0.05 ml per cm² of the leaf. After 60 minutes from the application, the plant 10 showed a surprisingly and non-expected improvement, in a more sur-

prisingly manner it was observed the total recovery of the leaf 11. This can be clearly appreciated in Figure 1B.

EXAMPLE 3

Application in plants obtained by micro-propagation

[0040] In order to study the effect of the composition of the present invention over plants obtained by micro-propagation, which goes from *in vitro* conditions to *ex vitro* conditions, two study groups of a species *Paulownia elongata* were formed. The composition prepared according to the example 1 was applied to the first group every 15 days, while the second group was not applied with the composition, i.e., it was a control group. After 45 days and three applications at 0, 15 and 30 days, the first group of plants showed a greater size than the second group of plants, further, showed great force and an intense green color as indicatives of having a good photosynthesis. All this was interpreted as a good adaptation of the plants to the *ex vitro* conditions. Contrariwise, the second group of plants showed a pale green or yellowish color, with less vegetative development, in other words, they showed less number of leaves and smaller size thereof compared to the leaves of the first group of plants.

EXAMPLE 4

Culture field studies of plants previously treated with the composition in the greenhouse

[0041] Some of the first group of plants, as well as others from the second group mentioned in the example 3 were transplanted to a culture field, wherein were treated under the same environment and watering conditions. After 6 months of said transplant, the plants of this group being treated with the composition of the present invention showed an important vegetative development and a normal development of the apical meristem of the plant, on the contrary, the plants not treated showed a smaller size and a non-efficient leaves development.

EXAMPLE 5

Application of the composition to greenhouse small leaf clusters

[0042] In order to determine the effect of the composition of the present invention applied to greenhouse tomato small leaf clusters of the "saladet" variety, 5 study groups were formed identified as follows:

- 1) C, a control group not receiving the composition of the present invention;
- 2) R1, a group to which the composition prepared according to the example 1 was applied, adding 3.0 ml/l of foliar fertilizer of the commercial trademark "Bayfolan Forte" and 3 mg of phloridzin (glucoside);

3) R2 a group to which the composition prepared according to the example 1 was applied, incorporating 3 mg of phloridzin in the above mentioned fertilizer;

4) R3 a group to which the composition prepared according to the example 1 was applied, i.e., without foliar fertilizer and without glucoside;

5) R4, a group to which the composition prepared according to the example 1 was applied, with 3ml/l of foliar fertilizer of the commercial trademark "Bayfolan Forte".

[0043] In groups R1 to R4 the composition was applied in an application dose of 0.05 ml per cm² of the leaf, such that all the upper surface of the leaf of the plant was covered.

[0044] The variable studied in the plants was the height in cm, after 3 applications in 45 days.

[0045] In an variance statistics analysis ($\alpha=0.05$) it was observed the existence of a significant difference between groups R1 to R4 with respect to the control group C, resulting in the conducting of a multiple media comparative analysis shown in Figure 3, wherein it is important to point out that averages with the same letter are not significantly different. In Figure 3, the existence of very important differences between groups R1 and R2 with respect to the control group C, to which the composition was not applied, such that in this development phase of the plants, the composition used in groups R1 and R2 favors a better growing of the plants. In order to make the above more evident, reference to Figure 2 is made, wherein a plant 20 of the control group C is shown compared to another plant belonging to the R1 group, wherein the height difference therebetween is evident.

EXAMPLE 6

Application to small leaf clusters transplanted to a culture field.

[0046] The same study groups as in example 5 were formed of the same age small leaf clusters, from the same tomato variety, but transplanted to soil, the response varieties were the plant height and stem diameter at the base thereof, said variables were measured after 4 applications and 65 days after the transplanting in the field.

[0047] Regarding the height variable, the variance analysis showed the existence of a significant difference ($\alpha=0.05$) in the different groups, leading to carry out a multiple average comparative test, shown in Figure 4, wherein it is important to mention that averages with the same letter are not significantly different. In Figure 4, it was observed a significant existence between the group of plants R2 which received the treatment and the control group, concluding that at least in this development phase, the plants of R2 treated with the formulation showed a bigger size with respect to the plants of the control group C.

[0048] Regarding the stem diameter variable, the variance analysis showed the existence of a significant difference ($\alpha=0.05$), also leading to carry out a multiple average comparative test shown in Figure 5, wherein it is important to mention that averages with the same letter are not significantly different. In Figure 5, it was observed a significant existence between the group of plants of R1 which received the treatment and the control group, concluding that at least in this development phase, the plants of R1 treated with the formulation showed a bigger diameter with respect to the plants of the control group.

Claims

1. An aqueous composition for the protection and strengthening of plants comprising:

- a) polyethylene glycol in a concentration from about 2.0 to about 10.0 grams per liter of the composition, wherein the polyethylene glycol has a molecular weight of about 6000 to about 8000 grams/mol;
- b) vitamin A or a derivative or precursor thereof, including mixtures of same, in a concentration of about 0.1 to about 10.0 grams per liter of the composition; and,
- c) at least a red coloring agent in a concentration of about 0.1 to about 10.0 grams per liter of the composition, wherein said coloring agent enables the passage of light at wavelengths greater than 600 nanometers

wherein the vitamin A derivative is selected from the group consisting of vitamin A acetate and vitamin A palmitate and wherein the vitamin A precursor is selected from the group consisting of beta-carotene, zeaxanthine and luteine.

2. The aqueous composition for the protection and strengthening of plants, according to claim 1, wherein the red coloring agent is a food grade coloring agent.

3. The aqueous composition for the protection and strengthening of plants, according to claim 2, wherein the red coloring agent is selected from the group comprising red 3 coloring agent, red 40 coloring agent or a mixture thereof.

4. The aqueous composition for the protection and strengthening of plants, according to claim 1, further comprising:

- d) at least a glucoside in a concentration from 0.5 to 15 mg per liter of the composition.

5. The aqueous composition for the protection and

strengthening of plants, according to claim 1, further comprising:

e) at least a foliar fertilizer

6. The aqueous composition for the protection and strengthening of plants according to claim 4, wherein said glucoside is selected from the group comprising phloridzin (phloretin-2- β -D-glucoside); phloretin (β -(4-hydroxyphenyl)-2,4,6-trihydroxypropiophenone), quercetin (3,3',4',5,6-pentahydroxyflavone), rutoside (quercetin-3-rutinoside) and phloroglucinol (1,3,5-trihydroxybenzene).

7. The aqueous composition for the protection and strengthening of plants according to claim 6, wherein said glucoside is phloridzin.

8. The aqueous composition for the protection and strengthening of plants according to claim 1, further having a pH from 3.5 to 7.0

9. An application method of the aqueous composition for the protection and strengthening of plants as defined in claim 1, said method comprising the steps of:

- a) providing a plant; and,
- b) applying the composition onto the leaves of the plant in an application dose of about 0.03 to about 0.05 ml per cm² of the leaf of the plant.

10. The application method of an aqueous composition for the protection and strengthening of plants, according to claim 9, wherein the application is carried out onto the upper side of the leaf of the plant.

11. The application method of an aqueous composition for the protection and strengthening of plants, according to claim 9, wherein the application is carried out by mechanical or manual spraying.

12. The application method of an aqueous composition for the protection and strengthening of plants, according to claim 9, wherein the composition is applied to plants obtained by micro-propagation going from *in vitro* conditions to *ex vitro* conditions.

13. The application method of an aqueous composition for the protection and strengthening of plants, according to claim 9, wherein the composition is applied to small leaf clusters obtained by seed germination and cuttings methods.

Patentansprüche

1. Eine wässrige Zusammensetzung zum Schutz und zur Verstärkung der Pflanzen, **dadurch gekennzeichnet**

zeichnet, dass sie enthält:

- a) Polyethylenglycol in einer Konzentration von 2.0 zu 10,0 g/L von der Zusammensetzung, wobei das Polyethylenglycol ein Molekulargewicht von ca. 6000 zu ca. 8000 g/mol hat;
- b) Vitamin A oder ein derivat oder ein Precursor davon, mitsamt deren Mischungen, in einer Konzentration von ca. 0.1 zu ca. 10,0 g/L von der Zusammensetzung hat; und,
- c) mindestens einem roten Farbstoff in einer Konzentration von ca. 0,1 zu ca. 10,0 g/L der Zusammensetzung hat, wobei dieser Farbstoff den Durchgang des Lichtes in einer Wellenlänge grösser als 600 nm erlaubt

wobei das Derivat des Vitamins A von einer Gruppe bestehend aus Vitamin A Azetat un Vitamin A Palmitat ausgewählt wird und wobei das Vitamin A Precursor von einer Gruppe bestehend aus β -Carotin, Zeaxanthin und Lutein ausgewählt wird.

2. Die wässrige Zusammensetzung zum Schutz und zur Verstärkung der Pflanzen nach dem Anspruch 1, wobei der rote Farbstoff lebensmitteltauglich ist.

3. Die wässrige Zusammensetzung zum Schutz und zur Verstärkung der Pflanzen nach dem Anspruch 2, wobei der rot Farbstoff von einer Gruppe bestehend aus Farbstoff Rot 3, Farbstoff Rot 40 oder eine Mischung von Beiden ausgewählt wird.

4. Die wässrige Zusammensetzung zum Schutz und zur Verstärkung der Pflanzen nach dem Anspruch 1, zusätzlich bestehend aus:

- d) mindestens einem Glucosid in einer Konzentration von 0,5 zu 15 mg/L von der Zusammensetzung; und/oder,

5. Die wässrige Zusammensetzung zum Schutz und zur Verstärkung der Pflanzen nach dem Anspruch 1, zusätzlich bestehend aus:

- e) mindestens einem Blattdüngemittel.

6. Die wässrige Zusammensetzung zum Schutz und zur Verstärkung der Pflanzen nach dem Anspruch 4, wobei das obgenannte Glucosid von Phlorizin (Phloretin-2- β -D-Glucosid); Phloretin (β -(4-Hydroxyphenyl)-2,4,6,-Trihydroxiropiophenon), Quercetin (3,3',4',5,6-Pentahydroxi flavon), Rutin (Quercetin-3-Rutinosid) oder Phloroglucinol (1,3,5-Trihydroxi-benzen) ausgewählt wird.

7. Die wässrige Zusammensetzung zum Schutz und zur Verstärkung der Pflanzen nach dem Anspruch 6, wobei das obgenannte Glucosid Phlorizin ist.

8. Die wässrige Zusammensetzung zum Schutz und zur Verstärkung der Pflanzen nach dem Anspruch 1 die weiterhin ein pH von 3,5 zu 7,0 hat
9. Eine Anwendungsverfahren der wässrigen Zusammensetzung zum Schutz und zur Verstärkung der Pflanzen nach dem Anspruch 1, wobei das Anwendungsverfahren aus den folgenden Schritte besteht:
- eine Pflanze versorgen; und,
 - diese Zusammensetzung in die Pflanzenblätter anlegen in einer Applikationsdosis von 0.03 zu 0,05 ml/cm² des Pflanzenblattes.
10. Das Anwendungsverfahren von der wässrigen Zusammensetzung zum Schutz und zur Verstärkung der Pflanzen nach dem Anspruch 9, wobei die Anwendung auf der oberen Seite jedes Pflanzenblattes durchgeführt wird.
11. Das Anwendungsverfahren von der wässrigen Zusammensetzung zum Schutz und zur Verstärkung der Pflanzen nach dem Anspruch 9, wobei die Applikation bei mechanischer oder manueller Besprühung durchgeführt wird.
12. Das Anwendungsverfahren von der wässrigen Zusammensetzung zum Schutz und zur Verstärkung der Pflanzen nach dem Anspruch 11, wobei die Zusammensetzung an Pflanzen angewandt wird, welche bei Mikrovermehrung erhalten werden und zugleich von *in vitro* zu *ex vitro* Bedingungen gewechselt haben.
13. Das Anwendungsverfahren von der wässrigen Zusammensetzung zum Schutz und zur Verstärkung der Pflanzen nach dem Anspruch 11, wobei die Zusammensetzung an kleine, durch Germination oder Vegetativer Vermehrung erhaltene Sämlinge angewandt wird.

Revendications

1. Une composition aqueuse pour la protection et le renforcement des plantes, caractérisée parce qu'elle comprend:
- polyéthylène glycol en une concentration de 2.0 a 10.0 grammes par litre de la composition; où le polyéthylène glycol à un poids moléculaire de 6000 a 8000 grammes/mole;
 - vitamine A ou un dérivé ou un précurseur de la même, contenant ses composants, en une concentration de 0.1 a 10.0 grammes par litre de la composition; et,
 - au moins un colorant rouge en une concentration de 0.1 a 10.0 grammes par litre de la com-

position où le colorant ici mentionné laisse passer la lumière en longueurs d'onde supérieures à 600 nanomètres

- où le dérivé de vitamine A est sélectionné du groupe qui consiste en acétate de vitamine A et palmitate de vitamine A et où le précurseur de vitamine A est sélectionné du groupe qui consiste en bêta carotène, zéaxanthine et lutéine.
2. La composition aqueuse pour la protection et le renforcement des plantes, en conformité avec la revendication 1, caractérisée aussi parce que le colorant rouge se sélectionne entre colorants de grade alimentaire.
3. La composition aqueuse pour la protection et le renforcement des plantes, en conformité avec la revendication 2, caractérisée aussi parce que le colorant rouge se sélectionne entre colorant rouge 3, colorant rouge 40 ou un mélange des deux.
4. La composition aqueuse pour la protection et le renforcement des plantes, en conformité avec la revendication 1, caractérisée aussi parce qu'elle comprend en plus:
- au moins un glucoside en une concentration de 0.5 a 15 mg par litre de la composition.
5. La composition aqueuse pour la protection et le renforcement des plantes, en conformité avec la revendication 1 caractérisée aussi parce qu'elle comprend aussi:
- au moins un fertilisant foliaire.
6. La composition aqueuse pour la protection et le renforcement des plantes, en conformité avec la revendication 4 caractérisée aussi parce que le glucoside mentionné se sélectionne entre phlorizine (phlorétine - 2-β-glucoside), phlorétine (β-4-hydroxyphényl)-2,4,6-trihydroxypropionophénone), quercétine (3,3', 4',5,6-pentahydroxyflavone), rutine (quercétine-3-rutinoside) ou phloroglucinol (1,3,5-trihydroxybenzène).
7. Une composition aqueuse pour la protection et le renforcement des plantes, en conformité avec la revendication 6, caractérisée aussi parce que le glucoside ici mentionné est phlorizine.
8. Une composition aqueuse pour la protection et le renforcement des plantes, en conformité avec la revendication 1 caractérisée aussi parce que qu'elle a un pH de 3.5 a 7.0
9. Une méthode d'application d'une composition

aqueuse pour la protection et le renforcement des plantes telle qu'elle est définie par la revendication 1, dite méthode étant caractérisée parce qu'elle comprend les suivantes étapes:

- a) fournir une plante; et
- b) appliquer la composition mentionnée sur les feuilles de la plante, en une dose d'application de 0.03 à 0.05 ml/cm² de la feuille de la plante.

5

10

10. La méthode d'application d'une composition aqueuse pour la protection et le renforcement des plantes en conformité avec la revendication 9, caractérisée aussi parce que l'application se réalise sur la face supérieure de chaque feuille de la plante.

15

11. La méthode d'application d'une composition aqueuse pour la protection et le renforcement des plantes en conformité avec la revendication 9, caractérisée aussi parce que l'application se réalise par aspersion mécanique ou manuelle.

20

12. La méthode d'application d'une composition aqueuse pour la protection et le renforcement des plantes en conformité avec la revendication 9, caractérisée aussi parce que la composition s'applique sur des plantes obtenues par micropropagation; ce qui passe d'être des conditions *in vitro* à des conditions *ex vitro*.

25

30

13. La méthode d'application d'une composition aqueuse pour la protection et le renforcement des plantes en conformité avec la revendication 9, caractérisée aussi parce que la composition s'applique à des plantules obtenues à travers des méthodes de germination de graines ou de boutures.

35

40

45

50

55




FIG. 1A




FIG. 1B




FIG. 2




FIG. 3




FIG. 4




FIG. 5

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 4094845 A [0009]
- WO 9624246 A1 [0010]
- WO 91003159 A [0011]
- US 4710394 A [0012]
- SU 1153403 A1 [0013]
- US 4828890 A [0014]