



Research article

Type C feruloyl esterase from *Aspergillus ochraceus*: A butanol specific biocatalyst for the synthesis of hydroxycinnamates in a ternary solvent system



Evelyn Romero-Borbón^a, Daniel Grajales-Hernández^b, Mariana Armendáriz-Ruiz^b, Lorena Ramírez-Velasco^b, Jorge Alberto Rodríguez-González^b, Luis Alberto Cira-Chávez^a, María Isabel Estrada-Alvarado^a, Juan Carlos Mateos-Díaz^{b,*}

^a Departamento de Biotecnología y Ciencias Alimentarias, Instituto Tecnológico de Sonora, 5 de Febrero 818 Sur, Centro, Ciudad Obregón, Sonora, Mexico

^b Departamento de Biotecnología Industrial, Centro de Investigación y Asistencia en Tecnología y Diseño del Estado de Jalisco, A.C. Camino Arenero 1227 El Bajío del Arenal, Zapopan, Jalisco, Mexico

ARTICLE INFO

Article history:

Received 12 February 2018

Accepted 27 June 2018

Available online 03 July 2018

Keywords:

Aspergillus ochraceus

Biocatalyst

Butanol specificity

Butanol

Butyl caffeate

Butyl hydroxycinnamates

Hydroxycinnamic acids

Hydroxycinnamoyl esterase

Solid state fermentation

Strain producer

Type C feruloyl esterase

ABSTRACT

Background: *Aspergillus ochraceus* was isolated from coffee pulp and selected as an interesting hydroxycinnamoyl esterase strain producer, using an activity microplate high-throughput screening method. In this work, we purified and characterized a new type C *A. ochraceus* feruloyl esterase (AocFaeC), which synthesized specifically butyl hydroxycinnamates in a ternary solvent system.

Results: AocFaeC was produced by solid state fermentation, reaching its maximal activity (1.1 U/g) after 48 h of culture. After purification, the monomeric protein (34 kDa) showed a specific activity of 57.9 U/mg towards methyl ferulate. AocFaeC biochemical characterization confirmed its identity as a type C feruloyl esterase and suggested the presence of a catalytic serine in the active site. Its maximum hydrolytic activity was achieved at 40°C and pH 6.5 and increased by 109 and 77% with Ca²⁺ and Mg²⁺, but decreased by 90 and 45% with Hg²⁺ and Cu²⁺, respectively. The initial butyl ferulate synthesis rate increased from 0.8 to 23.7 nmol/min after transesterification condition improvement, using an isoctane:butanol:water ternary solvent system, surprisingly the synthesis activity using other alcohols was negligible. At these conditions, the synthesis specific activities for butyl *p*-coumarate, sinapinate, ferulate, and caffeate were 87.3, 97.6, 168.2, and 234 U/μmol, respectively. Remarkably, AocFaeC showed 5 folds higher butyl caffeate synthesis rate compared to type B *Aspergillus niger* feruloyl esterase, a well-known enzyme for its elevated activity towards caffeic acid esters.

Conclusions: Type C feruloyl esterase from *A. ochraceus* is a butanol specific biocatalyst for the synthesis of hydroxycinnamates in a ternary solvent system.

How to cite: Romero E, Grajales D, Armendáriz M, et al. Type C feruloyl esterase from *Aspergillus ochraceus*: A butanol specific biocatalyst for the synthesis of hydroxycinnamates in a ternary solvent system. Electron J Biotechnol 2018;35. <https://doi.org/10.1016/j.ejbt.2018.06.004>.

© 2018 Pontificia Universidad Católica de Valparaíso. Production and hosting by Elsevier B.V. All rights reserved. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Feruloyl esterases (Faes; E.C. 3.1.1.73) are a group of hydrolases broadly distributed in plants and microorganisms that catalyze the hydrolysis or condensation between hydroxycinnamic acids (*p*-coumaric, caffeic, ferulic, and sinapinic acid) and an alcohol moiety. Their substrate specificity against four methyl hydroxycinnamates and their primary sequence identity allows the Faes to be classified into

four types (A–D) [1]. Type A Faes show a preference for methyl ferulate and sinapinate, type B Faes for methyl *p*-coumarate and caffeate, while type C and D Faes have a broad specificity, hydrolyzing all methyl hydroxycinnamates.

Nowadays, Faes from several fungi of the *Aspergillus* genus are produced by solid state fermentation (SSF) with several advantages, such as a high volumetric productivity. Furthermore, many agro-industrial by-products containing hydroxycinnamic acids, may be used as supports/substrates for the induction of Fae activity [2,3]. For example, wheat bran and sugar beet pulp are inducers for type A (AnFaeA) and type B (AnFaeB) feruloyl esterases from *Aspergillus niger*, respectively [4,5], while maize bran induces a type C feruloyl esterase from *Aspergillus terreus* [6].

* Corresponding author.

E-mail address: jcmateos@ciatej.mx (J.C. Mateos-Díaz).

Peer review under responsibility of Pontificia Universidad Católica de Valparaíso.

Faes have considerable roles in biotechnological processes in a wide variety of applications such as in biofuel, food, feed, pulp, paper, cosmetics, and pharmaceutical industries [7,8]. They have the ability to release ferulic acid and other hydroxycinnamic acids from plant biomass, which have been demonstrated to possess antioxidant, anti-inflammatory, antitumoral, antimicrobial and photoprotective activities [9] and are precursors of high economic value molecules [10,11,12]. In addition, when the acidic moiety of a hydroxycinnamic acid is esterified with an alcohol (aliphatic or sugar), its biological activities might be enhanced or modified. Furthermore, the hydrophilic–lipophilic balance modification of hydroxycinnamic acids after esterification may also allow its incorporation into water or oil based formulations, which could be of interest in the cosmetic, food, agro-, and pharmaceutical industries [9].

During the last 15 years, several research works demonstrated that alkyl hydroxycinnamates biological activities show a cut-off effect between C₄ and C₈ alkyl chain length [13,14,15]. In this respect, butyl and octyl esters of *p*-coumaric and caffeic acid, showed the highest *in vitro* and *in vivo* anti-inflammatory activity through NF- κ B pathway inhibition, suppressing the secretion of PGE2 [16]. Moreover, short and medium alkyl caffeate esters (<C₅) showed enhanced antitumoral activities against cervical and gastric cancer cells in comparison with the non-esterified acids [17], while the butyl ester of sinapinic acid is reported as a potent antioxidant inhibiting lipid oxidation [18]. Additionally, butyl esters of coumaric acid showed the highest antifungal activity against *Pyricularia grisea*, *Valsa mali* or *Botryosphaeria dothidea* [19]. Finally, it has been demonstrated that butyl ferulate is an important precursor of a novel prodrug, used as a lipophilic antioxidant [20]. All these previous works prove the biological importance of butyl hydroxycinnamates and the need of alkyl specific enzymes for their synthesis.

The derivatization of hydroxycinnamic acids with an alkyl alcohol has been performed by esterification or transesterification using Faes in a ternary solvent system [21,22,23,24]. This system contains a non-polar solvent, an alcohol (used as the substrate), and an aqueous phase containing the enzyme. It has been shown that the concentrations of the solvent, alcohol, and water in this reaction system strongly influence the Fae conversion yield [21,23,25]. The synthesis of hydroxycinnamic esters using Faes instead of chemical catalysts reduces the formation of secondary by-products, using environmentally friendly processes. Type C Faes possess a broad specificity during the synthesis of hydroxycinnamic acid esters. Hence, type C Faes from *Fusarium oxysporum* [26], *Myceliophthora thermophila* (formerly *Sporotrichum thermophile*) [25], and *Talaromyces stipitatus* [27] were successfully used in the synthesis of propyl, butyl and arabinosyl hydroxycinnamic acid esters reaching maximum yields of approximately 70% at mild temperatures (30–37°C). However, at our knowledge Faes alkyl specificity during butyl hydroxycinnamates synthesis has been poorly studied. Therefore, bioprospecting for alkyl specific Faes capable of synthesizing hydroxycinnamic butyl esters is still required.

In a previous work, *A. ochraceus* was selected among a fungal strain collection isolated from coffee waste and used to produce an enzymatic extract containing Fae activity, which released caffeic acid from coffee pulp [28,29]. In this work, a new type C feruloyl esterase from *A. ochraceus* (AocFaeC) was produced by SSF using maize bran as an inducer, purified, biochemically characterized and its biocatalytic potential was demonstrated for the alkyl specific synthesis of butyl esters of hydroxycinnamic acids, which own interesting biotechnological applications.

2. Materials and methods

2.1. Materials and strains

Bradford reagent, phenylmethanesulfonyl fluoride (PMSF), MgSO₄, K₂HPO₄, HCl, formic acid, urea, ammonium sulfate, hydroxycinnamic acids, methanol, ethanol, propanol, butanol, *tert*-butanol, octanol,

dodecanol, β - mercaptoethanol, bacteriological agar, bovine serum albumin (BSA), Tween 80, *n*-hexane, ethyl acetate, petroleum ether, diethyl ether, acetonitrile, buffer salts, and metal ion salts were purchased from Sigma (Toluca, Mexico). The broad range molecular weight standard was purchased from Bio-Rad (Ciudad de Mexico, Mexico). Maize bran was kindly provided by Minsa® (Los Mochis, Sinaloa, Mexico). *A. niger* strains for type A and type B feruloyl esterase production were kindly donated by Dr. Record and Dr. Lvasseur from INRA, France.

2.2. Fungal strain conservation and inoculum preparation

A. niger and *A. ochraceus* spores were maintained in 50% v/v glycerol at –80°C. The spore suspension used as inoculum was prepared in 250 ml Erlenmeyer flasks each containing 50 ml of the media reported by Record et al. [30] and Rodríguez et al. [31] for *A. niger* and *A. ochraceus*, respectively. For *A. ochraceus*, washed maize bran with a particle size of approximately 0.15 mm (40 g/l) was used as an inducer. Flasks were incubated for 4 d at 30°C, and spores were collected by stirring 50 ml of 0.01% w/v Tween 80 sterile solution.

2.3. Feruloyl esterase production in SSF

SSF was performed essentially as described by Rodríguez et al. [31]. Prior to fermentation, maize bran (substrate) was washed thoroughly with tap water to eliminate the soluble sugars. Then, it was dried at 30°C for 3 d and finally milled and sieved to achieve a particle size of approximately 0.42 to 0.59 mm. Polyurethane foam was cut into 0.5 cm cubes and employed as a support for the SSF. The impregnation media composition (g/l) was as follows: MgSO₄ (1); K₂HPO₄ (5); and urea (4). The support and substrate were mixed in a 1/3 (w/w) proportion, and then, each gram of mixture was added with 1.5-fold concentrated impregnation media at a pH of 6.5 and sterilized at 121°C for 15 min. The spore suspension was added to achieve 3 × 10⁷ spores per gram of dry matter (gdm). The humidity and temperature were adjusted to 75% w/w and 30°C, respectively. After culture, the solid ferment of each column was placed in a plastic bag. The enzymatic extract was obtained after mixing the solid ferment with 2.5 mM MOPS at a pH of 7.2 in a 1/2 (w/v) proportion and applying slight manual pressure to each bag. The enzymatic extract was centrifuged at 14,000 ×g for 5 min to eliminate fungal residues and suspended matter. Two fermentation columns were removed every 24 h to measure the Fae hydrolytic activity in the enzymatic extracts using methyl ferulate (MF) as the substrate and the spectrophotometric assay described below. Average values and standard deviation were reported.

2.4. Synthesis of methyl and butyl hydroxycinnamates

Each hydroxycinnamic acid was dissolved in methanol or butanol to obtain a 0.5 M solution, and then, 4% (v/v) HCl was added. The reaction mixture was heated under reflux for 18 h. The reaction progress was followed by thin layer chromatography (*n*-hexane/ethyl acetate; 2/1) and after complete reaction, the solvent was evaporated under reduced pressure using a rotary evaporator. The obtained powder was purified by silica gel column chromatography using different petroleum ether/diethyl ether mixtures. Fractions with the corresponding ester were pooled and dried to obtain white crystals with a purity of >98%.

2.5. Feruloyl esterase hydrolytic activity assays

2.5.1. Spectrophotometric assay

The Fae activity was determined by measuring the hydrolysis of different methyl hydroxycinnamates (5 mM) in the presence of the pH indicator 4-nitrophenol (0.5 mM), as described by Ramírez et al. [28]. The substrate solution was prepared using *tert*-butanol and 2.5 mM MOPS at a pH of 7.2 in a 1/9 (v/v) proportion. The decrease in

absorbance at 410 nm was continuously monitored in a microplate reader X-Mark from BioRad (Ciudad de Mexico, Mexico) at 30°C and a pH of 7.2 over 15 min. The reaction rate was calculated using the extinction coefficient obtained from standard curves (0 to 1 mM) for each hydroxycinnamic acid. Blank experiments were carried out using 2.5 mM MOPS buffer at a pH of 7.2. One unit corresponds to 1 μ mol of hydroxycinnamic acid released per minute.

2.5.2. Chromatographic assay

The Fae activity was also measured by quantifying the hydroxycinnamic acid released after 30 min of enzymatic hydrolysis of the corresponding ester in 50 mM potassium phosphate buffer at a pH of 7 and 30°C. Reaction products were monitored at 310 nm in a Varian HPLC System (California, USA) with a ZORBAX Eclipse XDB-C18 column (4.6 mm, 3 \times 250 mm, Agilent) at 30°C using water with 1% (v/v) formic acid and acetonitrile (60/40, v/v) as the mobile phase. The reaction rate was calculated using the extinction coefficient of each hydroxycinnamic acid, which was obtained from standard curves (0 to 5 mM). Blank experiments were carried out using 50 mM potassium phosphate buffer at a pH of 7. HPLC grade solvents were used during analysis. One unit corresponds to 1 μ mol of hydroxycinnamic acid released per minute.

2.6. Feruloyl esterase purification

Ammonium sulfate (2 M) was added to the enzymatic extract, which was filtered through a 0.45 μ m cellulose membrane (Millipore) and then loaded into a HiPrep Phenyl Sepharose FF 16/10 (Phenyl S) column from GE (Little Chalfont, United Kingdom) pre-equilibrated with 2.5 mM MOPS at a pH of 7.2 (buffer A) containing 2 M ammonium sulfate. The column was washed with 10 column volumes (CVs) of the same buffer to remove any unbound proteins. The elution was performed in 20 CVs using a linear gradient of 2–0 M ammonium sulfate in buffer A at a flow rate of 1 ml/min. Active fractions were pooled and dialyzed overnight against buffer A. Afterwards, the pooled fractions were loaded into a DEAE 16/10 from GE (Little Chalfont, United Kingdom) pre-equilibrated with buffer A, and then, the column was washed with 10 CVs using the same buffer. The protein elution was performed in 10 CVs with a linear gradient of 0–0.5 M NaCl in buffer A at a flow rate of 1 ml/min. Active fractions were pooled and concentrated to 200 μ l using an Amicon cellulose membrane from Millipore (10 kDa molecular mass cut-off) and loaded into a Superdex G200 16/60 from GE (Little Chalfont, United Kingdom) equilibrated with buffer A containing 150 mM NaCl. The elution was performed in 1.5 CVs, and active fractions were pooled and loaded onto a Mono Q 5/50 GL column from GE (Little Chalfont, United Kingdom) pre-equilibrated with buffer A. The purified protein was eluted in 10 CVs with a linear gradient of 0–0.5 M NaCl in buffer A at a flow rate of 1 ml/min. Recombinant type A and type B Faes from *A. niger* were produced and purified according to Record et al. [30] and Levasseur et al. [32], respectively.

2.7. Protein and electrophoresis assays

The protein concentration was determined using the Bradford microassay [33] and BSA as the reference protein. Sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) at 12% was performed to verify protein purity and to estimate the Fae molecular weight [34]. Zymography was carried out in 12% PAGE under non-denaturing conditions according to Camacho et al. [35] using MF as the substrate for the detection of the activity bands.

2.8. N-terminal amino acid sequence determination

The N-terminal sequence was determined by Edman degradation of an electroblotted pure deglycosylated protein sample (40 μ g) obtained from a polyvinylidene difluoride membrane using a Trans-Blot Turbo

Transfer Starter System from Bio-Rad (Ciudad de Mexico, Mexico). Analysis was carried out on a Microflex II instrument from Bruker (Massachusetts, USA).

2.9. Biochemical characterization of type C feruloyl esterase from *A. ochraceus*

2.9.1. Effects of the temperature and pH

The hydrolytic activity of the purified protein was evaluated at different temperatures (30 to 70°C) and pH values (4 to 10) using the chromatographic assay and MF as the substrate. For the pH assays, different solutions at 50 mM were used including the following: citrate buffer (pH = 4 to 5.5), sodium phosphate (pH = 6 to 7), Tris-HCl (pH = 7.5 to 8.5), and glycine-NaOH (pH = 9 to 10). Assay was performed by triplicate. Average values and standard deviation were reported.

2.9.2. Effects of metal ions and inhibitors

The effects of various metal ions (Ca^{2+} , Mg^{2+} , Mn^{2+} , Zn^{2+} , Cu^{2+} , and Hg^{2+}) and inhibitors (PMSF and β -mercaptoethanol) were evaluated by measuring the hydrolytic activity after 1 h of incubation using the chromatographic assay and MF as the substrate. The final concentration of all effectors was 1 mM. Assay was performed by triplicate. Average values and standard deviation were reported.

2.9.3. Kinetic parameters

The initial hydrolysis rates at different (0.01 to 4 mM) methyl caffeate (MC), methyl *p*-coumarate (MpC), MF, and methyl sinapinate (MS) concentrations were evaluated using the chromatographic assay. Afterwards, K_m and V_{max} were calculated using the Michaelis-Menten plot, Lineweaver-Burk regression plot and SigmaPlot 12.0 (Systat Software Inc.). For calculation of the k_{cat} , the molecular weight was estimated by SDS-PAGE. Assay was performed by triplicate.

2.10. Feruloyl esterase synthetic activity assays

2.10.1. Selection of the ternary solvent system for butyl ferulate synthesis

The synthesis of butyl ferulate was carried out by the transesterification of 50 mM MF in a ternary solvent system (isooctane:butanol:water) at 30°C, a pH of 7, and 1500 rpm in an Eppendorf Thermomixer[®]. The ternary solvent system was prepared according to Topakas et al. [21], with 1 ml being used in 1.5 ml sealed microtubes. To select butanol and water content in the ternary reaction system, seven isooctane: butanol: water relations (97.5:0.5:2; 92:6:2; 85.5:12.5:2; 73:25:2; 48:50:2; 74:25:1 and 71:25:4) were evaluated on the initial synthetic rate. The effect of enzyme concentration (1.3 to 20.7 μ g/ml), BSA concentration (0 and 1 mg/ml), and initial aqueous pH (3 to 9) was evaluated after 30 min of reaction on the initial synthetic rate in the ternary solvent system previously selected. For the pH assays, different solutions at 50 mM were used including the following: citrate buffer (pH = 4 to 5), MES (pH = 6 to 6.5), MOPS (pH = 7), Tris-HCl (pH = 8), and glycine-NaOH (pH = 9). Reaction samples (20 μ l) were taken at 0 as well as 30 min and stored at –20°C until HPLC analysis. Blank experiments were carried out without enzyme using the corresponding buffer. Assay was performed by triplicate. Average values and standard deviation were reported.

2.10.2. Synthesis of alkyl hydroxycinnamates

The transesterification of different methyl hydroxycinnamates (50 mM) with ethanol, propanol, butanol, octanol, and dodecanol was carried out in a ternary solvent system (isooctane:alcohol:water, 73:25:2, v:v:v) with 1 mg/ml BSA using 20 μ l of the purified enzyme (0.52 mg/ml) dissolved in 50 mM Tris-HCl buffer at pH 8. Reaction were performed in a 1 ml volume at 30°C and 1500 rpm for 4 h. Reaction samples (20 μ l) were removed at 0, 15, 30, 60, and 240 min, followed by storage at –20°C until HPLC analysis. Blank experiments were carried out using 50 mM Tris-HCl buffer at a pH of 8. Assay was




Fig. 1. Kinetics of *Aspergillus ochraceus* feruloyl esterase production by SSF. The production of feruloyl esterase is expressed in units (U) per gram of dry matter (gdm) using MF as a substrate.

performed by triplicate. Average values and standard deviation were reported.

2.10.3. Determination of feruloyl esterase synthetic activity by HPLC

Synthesis reaction samples (20 μ l) were diluted 10-fold with ethyl acetate and filtered through a 0.45 μ m filter before HPLC analysis. The reaction products were monitored at 310 nm in a Varian HPLC System (California, USA) using a Luna[®] 5 μ m silica (2) Phenomenex[®] column (100 \AA , 150 \times 4.6 mm, Agilent) at 30 $^{\circ}$ C using *n*-hexane/ethyl acetate (80/20) as the mobile phase. The reaction rate was calculated using the extinction coefficient of each butyl hydroxycinnamate, which was obtained from standard curve (0 to 1 mM). One unit corresponds to 1 μ mol of butyl hydroxycinnamate produced per minute. HPLC grade solvents were used during analysis. Assay was performed by triplicate. Average values and standard deviation were reported.

3. Results

3.1. Production and purification of type C feruloyl esterase from *A. ochraceus* (AocFaeC)

AocFaeC was produced by SSF using maize bran as an inducer (Fig. 1). Fae activity against MF was observed after 24 h, and the maximum activity of 1.1 U/gdm was reached around at 42 to 48 h of culture. Afterwards, the Fae activity decreased to 0.7 U/gdm after 120 h of fermentation. Thus, 200 g of solid ferment production was performed to accomplish the purification and characterization assays. The culture was stopped after 48 h to avoid possible protein proteolysis.

AocFaeC was purified using a four-step chromatographic strategy from 1.3 l of enzymatic extract (Table 1). For the first chromatography step, a hydrophobic Phenyl S column was used. This step allows 58-fold enrichment of the protein and the recovery of 109% of the total initial activity. In the second chromatography step, a weak anion exchange DEAE column was employed and allowed the protein to be enriched 80-fold with a 71.4% yield. Then, a molecular exclusion step

using a Superdex G200 column increased the purification factor by 295 with a 56.8% yield. Finally, a strong anion exchange Mono Q column led to the recovery of 0.23 mg of the pure protein (11.3% yield) with a purification factor of 483. The eluted protein was dialyzed overnight against 2.5 mM MOPS buffer at a pH of 7.2 to remove the residual NaCl. The homogeneity of the purified protein was verified by SDS-PAGE, Fig. 2a, showing a single band after purification. The specific activity of the pure AocFaeC was 57.9 U/mg using MF as a substrate (Table 1). The zymographic analysis through each purification step showed only one activity band when using MF as a substrate, Fig. 2b.

3.2. Biochemical characterization of AocFaeC

The molecular mass of AocFaeC was estimated to be approximately 34.6 kDa by SDS-PAGE, Fig. 2a, and 26.5 kDa (data not shown) by exclusion chromatography using a Superdex G200 column. The kinetic constants (K_m , V_{max} , and k_{cat}) were determined by measuring the initial hydrolysis rates of methyl hydroxycinnamates at different substrate concentrations (Table 2). The purified protein obeys Michaelis–Menten kinetics in the substrate concentration range of 0.01 to 4 mM. The K_m constants for all methyl hydroxycinnamates tested are shown in Table 2; for MS (1.11), the constant was approximately 2-fold higher than for MC (0.53), MpC (0.58), and MF (0.64). The V_{max} values against MC (40), MpC (47), MF (58) and MS (40) showed a broad hydrolytic capacity for *A. ochraceus* Fae. The substrate specificity was estimated by the specificity constant (k_{cat}/K_m). AocFaeC showed a slight preference for MF (31 356), MpC (28 037), and MC (26 113) with regards to MS (12 468). The AocFaeC N-terminal amino acid sequence is shown in Table 3. A BLAST analysis using the AocFaeC N-terminal sequence revealed more than 60% homology with other known N-terminal sequences of type C Fae from *Aspergillus* species and 100% homology with a putative type C Fae from *A. niger* (Table 3).

On the other hand, the activity of AocFaeC was measured as function of the temperature using MF as the substrate, Fig. 3a. AocFaeC showed a maximum activity at 40 $^{\circ}$ C, and then it decreased sharply to 7% of the maximum activity at 60 $^{\circ}$ C. The AocFaeC activation energy (E_a) was 24.9 kJ/mol, Fig. 3a. In the same way, the influence of the pH on the AocFaeC activity was measured in a pH range of 4 to 10, Fig. 3b. The maximum AocFaeC activity was observed at a pH of 6.5 using MF as a substrate, and surprisingly, the enzyme retained 40% of its maximum activity at a pH of 10.

The effects of various divalent metal ions and inhibitors on the AocFaeC hydrolytic activity using MF as the substrate are shown in Table 4. The AocFaeC activity increased with the addition of Ca^{2+} (209%), Mg^{2+} (117%), Mn^{2+} (143%), and Zn^{2+} (112%); was partially inhibited by β -mercaptoethanol (83%) and Cu^{2+} (55%); and a significant decrease in Fae activity was detected in the presence of Hg^{2+} (10%). Besides, the addition of PMSF resulted in a total loss of AocFaeC activity.

3.3. Effect of the ternary solvent system composition on the AocFaeC initial synthetic rate of butyl ferulate

The synthesis of butyl ferulate by the transesterification of MF with butanol was performed in an isoctane:butanol:water ternary system using AocFaeC at 30 $^{\circ}$ C (Fig. 4). The effects of the butanol, Fig. 4a, and

Table 1
Purification summary of *Aspergillus ochraceus* type C feruloyl esterase.

Step	Total activity (U)	Total protein (mg)	Specific activity (U/mg)	Yield (%)	Purification factor ^a
Crude extract	118.4	1029.3	0.12	100	1
Phenyl Sepharose	129.1	18.5	6.9	109	58
DEAE	84.6	8.7	9.6	71.4	80
Superdex G200	67.2	1.9	35.4	56.8	295
Mono Q	13.4	0.23	57.9	11.3	483

^a Purification factor = (Specific activity of each step) / (Specific activity of the crude extract).




Fig. 2. Electrophoretic analysis of the purified type C feruloyl esterase from *Aspergillus ochraceus*. (a) SDS-PAGE: lane 1, broad range molecular weight standards; lane 2, enzymatic crude extract; lane 3, pooled fractions eluted from the phenyl sepharose purification step; lane 4, pooled fractions eluted from the DEAE purification step; lane 5, pooled fractions eluted from the superdex G-200 purification step; and line 6, pooled fractions eluted from the mono Q purification step. The gel was stained with coomassie brilliant blue. (b) The Fae activity zymogram was determined using MF as the substrate.

water concentrations, Fig. 4b, as well as the pH in the aqueous phase, Fig. 4c, and enzyme concentration, Fig. 4d, were evaluated. The maximum initial synthetic rate of 2.8 nmol/min was reached at 25% (v/v) butanol, using an isooctane:butanol:water relation of 73:25:2 as seen in Fig. 4a. A higher butanol concentration using an isooctane:butanol:water relation of 48:50:2 sharply decreased the initial synthetic rate to 1.2 nmol/min (Fig. 4a). A lower (isooctane:butanol:water, 74:25:1 relation) or higher (isooctane:butanol:water, 71:25:4 relation) water concentration, decreased the initial synthetic rates to 0.9 nmol/min and 1.3 nmol/min, respectively (Fig. 4b). Thus, further experiments were carried out at an isooctane:butanol:water relation of 73:25:2. Afterwards, the addition of 1 mg/ml BSA into the aqueous phase of the ternary system reaction was evaluated. The addition of BSA increased the initial synthetic rate by 2.8-fold to 7.8 nmol/min. Regarding the effect of the initial aqueous phase pH on the initial butyl ferulate synthetic rate, AocFaeC was not active at an aqueous phase pH of 4, although the initial synthetic rate increased to 9.3 nmol/min at an aqueous phase pH of 9. It is important to mention that butyl ferulate was not detected using the corresponding buffer without enzyme, showing that alkaline catalysis is not occurring under the reaction conditions. Further studies were assessed at an aqueous phase pH of 8 to avoid possible methyl hydroxycinnamate degradation. Finally, the effect of the AocFaeC concentration on the initial synthetic rate was evaluated, Fig. 4d. The initial synthetic rate increased linearly to reach its maximum of 23.7 nmol/min using 10 μ g of AocFaeC. A higher enzyme concentration (20 μ g) did not increase the initial synthetic rate (23 nmol/min). Thus, further studies were performed using a ternary solvent composition of isooctane:butanol:water 73:25:2, an AocFaeC concentration of 10 μ g, 1 mg/ml BSA and an initial aqueous phase pH of 8, which increased the initial transesterification synthetic rate for obtaining butyl ferulate up to 30-fold (from 0.8 nmol/min to 23.7 nmol/min).

Table 2
Kinetic parameters of AocFaeC for different methyl hydroxycinnamates.

Substrate	K_m (mM) ^a	V_{max} (μ mol/min)/mg ^a	k_{cat} (1/min)	k_{cat}/K_m (1/mM min)
MC	0.53	40	13,840	26,113
MpC	0.58	47	16,262	28,037
MF	0.64	58	20,068	31,356
MS	1.11	40	13,840	12,468

^a Kinetic parameters were determined using methyl caffeate (MC), methyl *p*-coumarate (MpC), methyl ferulate (MF), and methyl sinapinate (MS) as substrates and determined from Lineweaver–Burk plots.

Table 3
N-terminal amino acid sequences of type C Fae from the *Aspergillus* genus.

Type C Fae source	N-terminal amino acid sequences	NCBI reference sequence
<i>A. ochraceus</i>	ANSPGCGKNP	
<i>A. niger</i>	ANSPGCGKNP	XP_001395336.1
<i>A. oryzae</i>	ANSSGCGKQP	XP_001820736.2
<i>A. udagawae</i>	TRSSGCGKQP	GAO87330.1
<i>A. lentulus</i>	TRSSGCGKQP	GAQ06322.1

3.4. Potential of AocFaeC for the synthesis of alkyl hydroxycinnamates

The transesterification of methyl to alkyl ferulates in the ternary solvent system conditions previously selected were performed using pure preparations of AnFaeA, AnFaeB, and AocFaeC as biocatalysts, in order to evaluate its synthetic potential. The specific synthetic activity per μ mol of pure enzyme for each Fae is shown in Fig. 5. The specific synthetic activities of AnFaeA for ethyl, propyl, butyl, octyl, and dodecyl ferulate were 19.0, 33.4, 20.5, 4.5, and 3.2 U/ μ mol of enzyme, respectively, showing a preference for the propyl chain. Aversely, AnFaeB shows low specific synthetic activity for all alkyl ferulate (around 2.9–5.4 U/ μ mol), without showing any alkyl synthesis preference. Interestingly, AocFaeC was able to synthesize only butyl and octyl ferulate with a specific synthetic activity of 168.2 and




Fig. 3. Effects of temperature and pH on the hydrolytic activity of AocFaeC. The relative activity was measured at different temperatures (a) and pH values using citrate buffer, sodium phosphate, Tris–HCl, and glycine–NaOH (b). The activation energy (E_a) was calculated using the Arrhenius equation. The activity was measured using MF as the substrate.

Table 4
Effects of metal ions and inhibitors on the AocFaeC hydrolytic activity.

Substance	Relative activity (%)
Control	100
CaCl ₂	209 ± 20
MgCl ₂	177 ± 4
MnCl ₂	143 ± 23
ZnCl ₂	112 ± 8
CuCl ₂	55 ± 4
HgCl ₂	10 ± 1
β-mercaptoethanol	83 ± 3
PMSF	0

5.3 U/μmol, respectively. As the specific synthetic activity of AocFaeC using other alcohols during transesterification reaction was negligible, it might be considered as a butyl chain specific Fae.

Finally, the transesterification of methyl to butyl hydroxycinnamates using AnFaeA, AnFaeB, and AocFaeC were performed to evaluate the acyl substituent preference (Fig. 6). The specific synthetic activities of AnFaeA for butyl *p*-coumarate, ferulate and sinapinate transesterification were 4.1, 20.5, and 39.3 U/μmol, respectively, showing a preference for butyl sinapinate synthesis. The specific synthetic activities of AnFaeB for butyl ferulate, caffeate, and *p*-coumarate transesterification were 5.4, 47.7, and 80 U/μmol, respectively, showing a preference for the synthesis of hydroxylated hydroxycinnamates. Interestingly, the specific synthetic activities of AocFaeC for the transesterification of butyl *p*-coumarate, sinapinate, ferulate, and caffeate were 87.3, 97.6, 168.2, and 234 U/μmol, respectively, showing the highest activity for the synthesis of butyl caffeate. AocFaeC shows 2.5- and 5-fold higher specific synthetic activities than AnFaeA and AnFaeB for butyl sinapinate and caffeate, respectively. These results demonstrate that at the conditions described in this work, AocFaeC was able to specifically synthesize all butyl hydroxycinnamates tested, at higher rates than AnFaeA and AnFaeB.

4. Discussion

After the first purification step, 109% of the total initial activity was recovered probably due to the presence of an inhibitor in the crude extract. AocFaeC is a monomeric protein and a type C feruloyl esterase like Fae from *Aspergillus oryzae* (AoFaeC) [36], *Talaromyces stipitatus* (TsFaeC) [37], *Myceliophthora thermophila* (StFaeC) [38], and *Fusarium oxysporum* (FoFaeC) [39]. The molecular weight for AocFaeC estimated by SDS-PAGE (34.6 kDa) was between those of AtFae-2 (23 kDa) and AtFae-3 (36 kDa) from *A. terreus* [8] and between those of StFaeC (23 kDa) [38], TsFaeC (55.3 kDa) [37], FoFaeC (62 kDa) [39], and AoFaeC (75 kDa) [40]. Thus, the type C feruloyl esterase group is composed of a wide range of molecular weight proteins and AocFaeC belongs to the medium molecular weight ones. The AocFaeC specific hydrolysis activity against MF (57.9 U/mg) was similar to the activity of TsFaeC (68.5 U/mg) [37] and significantly greater than that of StFaeC (0.7 U/mg) [25].

The comparison of k_{cat}/K_m values showed the broad specificity of AocFaeC against methyl hydroxycinnamates. Presently, the other reported fungi Fae that exhibit such a broad activity spectrum towards hydroxycinnamic esters are AtFae-2 [8], AtFae-3 [8], FoFae-I [21], FoFaeC [39], TsFaeC [37], AoFaeC [40], and StFaeC [25]. Comparing different type C Fae, the AocFaeC specificity constant using MF as a substrate is 3- and 330-fold higher than those of AoFaeC [40] and StFaeC [25], respectively, but is half of that reported for FoFaeC [39]. Nonetheless when comparing AocFaeC with type A Fae, the specificity constant using MF as a substrate is 6- and 18-fold greater than those values reported by Faulds et al. [41] and Schär et al. [42] for AnFaeA, which is an enzyme evolutionarily adapted to hydrolyze hydroxycinnamic acid esters containing methoxy substituents in the phenolic ring [43]. Thus, the broad substrate specificity and the N-terminal amino acid sequence reveals that the Fae from *A. ochraceus* corresponds to a type C feruloyl esterase [3].

The maximum activity of AocFaeC was reached at 40°C, similar to AtFae-2 and AtFae-3 from *A. terreus* [8]. However, this temperature




Fig. 4. Influence of the ternary solvent system composition on the AocFaeC initial transesterification synthetic rate of MF to BF. (a) Effect of different butanol concentrations. (b) Effect of different water concentrations. (c) Effect of the initial aqueous phase pH: citrate buffer, MES, MOPS, Tris-HCl, and glycine-NaOH. (d) Effect of different enzyme concentrations. All reactions were performed at 30°C.




Fig. 5. Specific activities of different feruloyl esterases in the synthesis of alkyl ferulates by transesterification. Comparison of the specific synthetic activity per μmol of pure preparations of feruloyl esterase type A from *Aspergillus niger* (AnFaeA), type B from *Aspergillus niger* (AnFaeB) and type C from *Aspergillus ochraceus* (AocFaeC). The transesterification reactions were performed using MF as substrates and a ternary solvent system isooctane:alcohol:water, 73:25:2 containing 1 mg/ml of BSA at a pH of 8 and 30°C.

differs from those reported for other mesophilic fungi type C Faes such as *Fusarium oxysporum* (65°C) [39] and *Talaromyces stipitatus* (60°C) [37]. On the other hand, the activation energy (E_a) of AocFaeC (24.9 kJ/mol) is similar to that of the type A Fae from *Aspergillus awamori* (21.4 kJ/mol) [44]. The effect of pH on AocFaeC hydrolytic activity towards MF reveals that the maximum activity was reached at a pH of 6.5, a value slightly higher than those reported for mesophilic fungi type C Faes, which reported pH values of 5 or 6 [8,37,39,40]. As shown in Fig. 3b, AocFaeC retains more than 60% of its maximum activity at pH values from 5.5 to 9.0, a property that might be exploited in alkaline processes.

In the effectors study, most metal ions tested increased the AocFaeC hydrolytic activity towards MF. The activation of AocFaeC hydrolytic activity by Ca^{2+} using MF as a substrate (209%), was higher than the activation of *Penicillium pinophilum* Fae using the same substrate (173%) [45] and that found for AwFaeA using sugarcane bagasse as a substrate (170%) [44]. Similar behaviors by Mg^{2+} and Mn^{2+} have been reported for the *Penicillium pinophilum* Fae [45] and a soil metagenomic library Tan410 Fae [46], respectively. The AocFaeC




Fig. 6. Specific activities of different feruloyl esterases in the synthesis of butyl hydroxycinnamates by transesterification. Comparison of the specific synthetic activity per μmol of pure preparations of feruloyl esterase type A from *Aspergillus niger* (AnFaeA), type B from *Aspergillus niger* (AnFaeB) and type C from *Aspergillus ochraceus* (AocFaeC). The transesterification reactions were performed using MC, MpC, MF, and MS as substrates and a ternary solvent system with the following composition: isooctane:butanol:water, 73:25:2 containing 1 mg/ml of BSA at a pH of 8 and 30°C.

hydrolytic activity increased to 112% after incubation with Zn^{2+} , unlike *Russula virescens* Fae, whose hydrolytic activity measured with MF was reduced to 98% [47]. Heavy metals such as Hg^{2+} and Cu^{2+} are thiol group and disulfide bond reducing agents [48], which inhibited AocFaeC activity by 90 and 45%, respectively. As AocFaeC maintains 83% of its activity after incubation with β -mercaptoethanol, a disulfide bond reducing agent [49], it can be suggested that thiol groups may play an important role in Fae activity. The complete loss of AocFaeC activity by PMSF reveals the presence of a serine in the active site [50].

In the transesterification of MF to butyl ferulate using the best ternary solvent system conditions found in this work (isooctane:butanol:water, 73:25:2, v:v:v), the AocFaeC initial synthetic rate (23.7 nmol/min or 2.8% conversion/h) was 16- and 35-fold higher than those of StFaeC (0.18% conversion/h) [25] and FoFae-I (0.08% conversion/h) [21] using hexane as a co-solvent, 43.4 and 50.8% of butanol, and 3.2 and 2% of water, respectively. In our case, reducing the butanol concentration in the ternary solvent system causes a loss of the microemulsion stability. Interestingly, AocFaeC initial synthesis rate increases with pH, as reported for FaeA1 and MtFae1a expressed in *Myceliophthora thermophila*, whose maximal hydrolytic activity was reached at pH 7 [51,52], while optimal synthesis activity was found at pH 8 [53]. This could probably be explained by the different ionization state of AocFaeC at the evaluated conditions, as stated by Antonopoulou et al. [53]. As in hydrolysis, AocFaeC shows a broad synthetic activity spectrum towards hydroxycinnamic methyl esters. However, differences between the hydrolytic and synthetic activity patterns were observed. AocFaeC preferentially hydrolyzes MF, while in transesterification reactions, it preferentially uses MC as a substrate, similar to as found for FoFae-I [21] and StFaeC [25]. Hence, this preference switch was probably caused by the ternary solvent system composition [54]. Nowadays, Faes alkyl chain specificity during transesterification of methyl hydroxycinnamates has been poorly studied. Schär and Nyström [55] evaluated the alkyl chain specificity of *Rhizomucor miehei* lipase in the synthesis of alkyl ferulates by direct esterification, proving its short alkyl chain preference (C_2 - C_4). To the best of our knowledge, AocFaeC is the only esterase that may perform the specific synthesis of butyl ferulate by transesterification of methyl ferulate. Finally, the AocFaeC synthetic specific activity per μmol of biocatalyst in the transesterifications of MF and MC to butyl ferulate and butyl caffeate was higher than that of AnFaeA and AnFaeB, which are naturally adapted for catalyzing methyl hydroxycinnamates with methoxylated or hydroxylated substituents in the aromatic ring, respectively. In this study, it was demonstrated that *Aspergillus ochraceus* type C Fae possesses a broad specificity for hydrolyzing methyl hydroxycinnamates and has the potential of carrying out the specific synthesis of butyl hydroxycinnamates, particularly by butyl caffeate, which possess all interesting antioxidant and anticarcinogenic properties [56].

Declarations of interest

None.

Acknowledgements

Evelyn Romero and Daniel Grajales acknowledge the scholarships received from Consejo Nacional de Ciencia y Tecnología. The authors acknowledged Dra. Régine Lebruns for determining the N-terminal amino acid sequence and Dr. Eric Record and Dr. Anthony Levasseur for the kind donation of *Aspergillus niger* transformants (CNRS, Marseille, France).

Financial support

This work was supported by the Fondo Mixto del Gobierno del Estado de Jalisco [project number 2012-07-190698].

- [53] Antonopoulou I, Leonov L, Jütten P, et al. Optimized synthesis of novel prenyl ferulate performed by feruloyl esterases from *Myceliophthora thermophila* in microemulsions. *Appl Microbiol Biotechnol* 2017;101(8):3213–26. <https://doi.org/10.1007/s00253-017-8631-8>.
- [54] Klibanov MA. Enzymatic catalysis in anhydrous organic solvents. *Trends Biochem Sci* 1989;14(4):141–4. [https://doi.org/10.1016/0968-0004\(89\)90146-1](https://doi.org/10.1016/0968-0004(89)90146-1).
- [55] Schär A, Nyström L. High yielding and direct enzymatic lipophilization of ferulic acid using lipase from *Rhizomucor miehei*. *J Mol Catal B: Enzym* 2015;118:29–35. <https://doi.org/10.1016/j.molcatb.2015.04.011>.
- [56] Serafim TL, Carvalho FS, Marques MPM, et al. Lipophilic caffeic and ferulic acid derivatives presenting cytotoxicity against human breast cancer cells. *Chem Res Toxicol* 2011;24(5):763–74. <https://doi.org/10.1021/tx200126r>.